

# GLYCOSYLATED HAEMOGLOBINS IN WOMEN WITH LOW RISK FOR DIABETES IN PREGNANCY

D F M Loke, L P Kek, S Chua, A C Thai, S S Ratnam

## ABSTRACT

*Glycosylated haemoglobin levels (HbA<sub>1c</sub>) were determined in 489 normal pregnant Singaporean women, attending a diabetic screening programme using a 50g glucose challenge test. All subjects had no risk factors for diabetes mellitus nor a history of previous gestational diabetes. They were selected when the 1h 50g glucose challenge (GCT) is < 6.66 mmol/l or a 2h post-glucose level of < 7.77 mmol/l (a 75g OGTT is repeated within a week of an abnormal GCT). Another 18 subjects with normal OGTT but did not have a 50g glucose challenge done were also included in the study. Mean and normal range (2 SD) of HbA<sub>1c</sub> levels in all subjects were 4.64% and 3.66-5.62%, respectively. Mean HbA<sub>1c</sub> levels also varied with gestation and ethnic group. The HbA<sub>1c</sub> appeared to be low at early gestation with nadir at 21-24 weeks and thereafter returned to initial levels at term. In Chinese, mean HbA<sub>1c</sub> levels, random plasma glucose and 1h glucose challenge were significantly higher than those of the Malays but not the Indians. In referring to these levels, the variation within gestation and ethnic group must also be considered.*

**Keywords:** glycosylated haemoglobin, pregnancy

SINGAPORE MED J 1995, Vol 36: 501-504

## INTRODUCTION

Glycosylated haemoglobin (HbA<sub>1c</sub>) measurements have helped in the assessment of treatment of diabetic patients<sup>(1-3)</sup>. They have proved valuable as an additional parameter for follow-up of pregnant diabetics where optimal control is of importance. While HbA<sub>1c</sub> levels have been determined in various pregnant populations, there is no standard reference range for comparison between populations. This is because different methods are used for measurement and also because standard reference materials are lacking.

In an attempt to define a reference range for our local population, we measured the HbA<sub>1c</sub> levels of a group of normal pregnant women at different gestation periods undergoing a screening programme for diabetes mellitus using a 50g glucose challenge test (GCT)<sup>(4)</sup>. This paper reports the HbA<sub>1c</sub> levels in women with a normal glucose challenge alone or with an abnormal GCT but a normal oral glucose tolerance test (75g OGTT) and we also compare the levels in different ethnic groups and gestation periods.

---

Department of Obstetric & Gynaecology  
National University of Singapore  
National University Hospital  
Kent Ridge Road  
Singapore 119074

D F M Loke, MSc  
Research Fellow

S Chua, MRCOG  
Senior Lecturer

S S Ratnam, FRCOG  
Professor & Head

Department of Medicine  
National University of Singapore

A C Thai, M Med (Int Med)  
Associate Professor

Gynecology Associates Pte Ltd  
#12-18 Mt Elizabeth Medical Centre  
Singapore 228510

L P Kek, MRCOG  
Consultant

Correspondence to: Dr D F M Loke

---

## MATERIALS AND METHODS

Subjects were recruited from the ante-natal clinic at the National University Hospital. These women did not have any historical or obstetrical risk factors for diabetes mellitus. On admission into the study, 2 venous blood samples (random samples) were drawn for glucose (KF coated tube) and HbA<sub>1c</sub> (EDTA-coated tube) before a 50g glucose load was given to the patient. Another venous blood sample was collected one hour after the glucose load for glucose estimation. Plasma glucose was determined by the hexo-kinase method<sup>(5)</sup> on the Cobas Mira. Inter-assay precision was 2%. Samples for HbA<sub>1c</sub> were haemolysed on the day of collection and stored at 4°C and analysed within 4 days by affinity chromatography<sup>(6)</sup> using the Glycogel (Pierce) Test Kit. Inter-assay precision was 5% at 5.3% HbA<sub>1c</sub> level and 4% at 16% HbA<sub>1c</sub> level.

Patients with a 1h glucose challenge  $\geq 6.66$  mmol/l were recalled within a week for a 75g glucose tolerance test. Only subjects with a normal GCT (1h glucose challenge  $< 6.66$  mmol/l) or an abnormal GCT but normal OGTT (2h post-glucose level  $\leq 7.77$  mmol/l by WHO criteria<sup>(7)</sup>) were included in the study. An additional 18 subjects who were not screened by a glucose challenge but had a normal 75g OGTT response were also included in the study.

To investigate the variation of HbA<sub>1c</sub> with duration of gestation, the subjects were grouped in smaller time intervals (Table I). As there were less subjects at early (1-12 weeks) and late (37-42 weeks) gestation, the time interval for these periods were longer to include a substantial number of subjects for each interval.

Laboratory values are reported as mean  $\pm 1$  standard deviation of the mean. Comparison of gestation periods between different ethnic groups was done by analysis of variance and Duncan test. To adjust for differences in gestation between ethnic groups, the data were compared by analysis of covariance with gestation as covariate. Similarly, comparison of data between different gestation groups was done by analysis of covariance with race as covariate. The data were also submitted to stepwise multiple regression analysis until all independent factors were significant at  $p < 0.05$  level. The statistical package, SAS, was used for analysis.

## RESULTS

Table II shows the mean HbA<sub>1c</sub> and glucose levels in normal pregnant women. Mean and range (2 SD limits) of HbA<sub>1c</sub> levels

were 4.64% and 3.66% to 5.62% respectively.

When subjects with normal glucose response were analysed according to ethnic groups, the mean gestation of the Malays was significantly higher than that of the Chinese or Indians (Table III). Before adjustment for differences in gestation, there was no significant differences in mean HbA<sub>1</sub> levels between different ethnic groups. However, after adjustment for differences in gestation, the Chinese were found to have significantly higher mean HbA<sub>1</sub> levels than the Malays (Table III). There were no significant differences in fasting plasma glucose (0h GTT) and glucose tolerance test levels between all groups. However, random plasma glucose and 1h post-glucose challenge levels were significantly higher in the Chinese than the Malays or Indians (p<0.01).

**Table I - Adjusted\* mean HbA<sub>1</sub> (±SD) levels in low risk pregnant women at various gestation.**

Gestation Group No	Weeks	No of Subjects	HbA <sub>1</sub> (%) Normal Group
1	1-12	19	4.74 ± 0.48
2	13-16	74	4.67 ± 0.48
3	17-20	97	4.62 ± 0.48
4	21-24	91	4.45 ± 0.48 <sup>a</sup>
5	25-28	63	4.65 ± 0.48
6	29-32	61	4.75 ± 0.47
7	33-36	70	4.76 ± 0.48
8	37-42	14	4.76 ± 0.48

\*Means adjusted for differences in race  
<sup>a</sup>p<0.05 vs all Gestation Groups.

**Table II - Mean (±SD) glucose and HbA<sub>1</sub> levels in low risk pregnant women with normal glucose response.**

Parameter	Mean ± SD	No.
Gestation (weeks)	23.7 ± 7.5	489
Random plasma glucose (mmol/l)	5.30 ± 1.38	471
1h glucose challenge (mmol/l)	6.75 ± 1.38	471
0h GTT (mmol/l)	4.29 ± 0.39	219
1h GTT (mmol/l)	7.46 ± 1.42	219
2h GTT (mmol/l)	6.10 ± 1.01	220
HbA <sub>1</sub> (%)	4.64 ± 0.49	489

**Table III - Adjusted\* mean (±SD) glucose and HbA<sub>1</sub> levels in low risk pregnant women of different races with normal glucose response.**


Parameter	Chinese	Malays	Indians
Gestation (weeks)	23.1 ± 7.2 (306)	25.6 ± 8.0 (137) <sup>a</sup>	21.9 ± 7.2 (46)
Random plasma glucose (mmol/l)	5.46 ± 1.00 (295) <sup>b</sup>	5.03 ± 1.01 (132)	5.02 ± 1.00 (44)
1h glucose challenge (mmol/l)	6.78 ± 1.36 (295) <sup>c</sup>	6.28 ± 1.38 (132)	6.02 ± 1.36 (44)
0h GTT (mmol/l)	4.29 ± 0.39 (158)	4.28 ± 0.39 (48)	4.27 ± 0.39 (13)
1h GTT (mmol/l)	7.42 ± 1.38 (158)	7.66 ± 1.38 (47)	9.28 ± 1.38 (13)
2h GTT (mmol/l)	6.15 ± 0.99 (158)	6.05 ± 0.99 (48)	5.77 ± 0.99 (13)
HbA <sub>1</sub> (%)	4.68 ± 0.48 (306) <sup>d</sup>	4.57 ± 0.49 (137)	4.60 ± 0.48 (46)

\*Means adjusted for differences in gestation. Numbers in brackets denote number of results for the particular parameter. <sup>a</sup>p<0.05 vs Chinese or Indians; <sup>b</sup>p<0.01 vs Indians and p<0.001 vs Malays; <sup>c</sup>p<0.001 vs Malays or Indians; <sup>d</sup>p<0.05 vs Malays.

Mean HbA<sub>1</sub> levels decreased from early gestation to a minimum at 21-24 weeks and increased to initial levels at term (Table I). Mean HbA<sub>1</sub> level at 21-24 weeks was significantly lower (p<0.05) than levels at other gestation periods.


Fig 1 shows the trend in mean HbA<sub>1</sub>, random plasma glucose and 1h post-glucose challenge levels in all subjects (471) with a glucose challenge done at various gestation periods. Mean random plasma glucose level was significantly lower at 17-20 weeks than 25-28 weeks (p<0.05) while 1h post-glucose challenge levels at 33-36 weeks were significantly lower than those at 13-16 or 29-32 weeks (p<0.05).

**Fig 1 - Mean (±SD) HbA<sub>1</sub>, mean random plasma glucose (RPG) and 1h glucose challenge (1h GC) levels in low risk pregnant women with normal glucose response.**



Footnote:  
n: number of subjects per gestation group; a:p<0.05 vs all groups; b:p<0.05 vs 13-16 & 29-32 weeks gestation groups; c:p<0.05 vs 25-28 weeks gestation group.

**Fig 2 - Mean (±SD) HbA<sub>1</sub>, mean fasting plasma glucose (0h), 1h and 2h post glucose loading (1h and 2h GTT) levels in low risk pregnant women with normal glucose response.**



Footnote:  
n: number of subjects per gestation group;  
a:p<0.05 vs 13-16, 29-32 & 33-36 weeks gestation groups;  
b:p<0.05 vs 17-20 & 21-24 weeks gestation groups;  
c:p<0.05 vs 1-12, 13-16 & 37-42 weeks gestation groups;  
d:p<0.05 vs 1-12 to 21-24 & 37-42 weeks gestation groups;  
e:p<0.05 vs 1-12 to 21-24 weeks gestation groups;  
f:p<0.05 vs 1-12 weeks gestation group;  
g:p<0.05 vs 1-12, 13-16, 25-28 & 33-36 weeks gestation groups;  
h:p<0.05 vs 1-12 & 13-16 weeks gestation groups.

Fig 2 shows the trend in mean HbA<sub>1c</sub>, 0h, 1h and 2h GTT levels in subjects (221) with a GTT done at various gestation periods. Mean HbA<sub>1c</sub> at 21-24 weeks was significantly lower than levels at 13-16, 29-32 and 33-36 weeks gestation. Mean fasting glucose levels, 0h GTT, at 17-20 and 21-24 weeks were significantly lower than levels at 1-12, 13-16, 25-28 and 33-36 weeks gestation. Mean 1h GTT levels at 25-28 and 29-32 weeks were significantly higher than those at 1-12 to 21-24 weeks gestation periods. Similarly, mean 2h GTT level at 25-28 weeks was significantly higher than those at 1-12, 13-16 and 37-42 weeks gestation periods and that at 29-32 weeks was significantly higher than those at 1-12 to 21-24 and 37-42 weeks gestation periods.

The results of multiple regression analysis of HbA<sub>1c</sub> with gestation, race, random plasma glucose and 1h glucose challenge are given in Table IV. Only gestation and 1h glucose challenge were significantly correlated with HbA<sub>1c</sub>.

## DISCUSSION

The mean and range of HbA<sub>1c</sub> levels in this study group with normal 1h glucose challenge or a normal 75g OGTT were 4.64% and 3.66%-5.62%, respectively. In an earlier study<sup>(8)</sup>, we used an electro-phoretic method to measure the HbA<sub>1c</sub> levels in a smaller group of 33 pregnant women with normal 50g OGTT response. These subjects were different from the low risk patients in our current study in that they had risk factors for diabetes mellitus. The mean and normal range of HbA<sub>1c</sub> levels then were 6.34% and 4.96% to 8.11% respectively. The reason for higher levels of HbA<sub>1c</sub> obtained in the previous study is probably due to difference in the method of estimation of HbA<sub>1c</sub>.

**Table IV - Multiple regression analysis of HbA<sub>1c</sub> with gestation, race, random plasma glucose and 1h glucose challenge**

Variable	b	F	p value	r
Intercept	4.056	999.3	0.0001	
1h glucose challenge	0.006	17.3	0.0001	0.18
Gestation	0.004	4.8	0.0292	0.10
			R	0.21

r, R - partial and multiple correlation coefficient. Only significant partial correlation coefficients (p<0.05) obtained after elimination by stepwise multiple regression analysis are listed.

HbA<sub>1c</sub> levels in our population of pregnant women were found to vary with ethnic groups. The Chinese had significantly higher levels of HbA<sub>1c</sub>, random plasma glucose and 1h post-glucose challenge glucose levels than the Malays. These differences could not be due to difference in gestation of the 2 groups since this variation had been taken into consideration in the analysis of the results. While 1h post-glucose challenge was observed to be significantly correlated to HbA<sub>1c</sub> levels, the variation of HbA<sub>1c</sub> cannot be satisfactorily explained by the variation in 1h glucose challenge since this factor only contributes about 3% of the variation (r=0.18, r<sup>2</sup>=0.03). There maybe other factors including socio-economic status, weight for height gain, and dietary habits which affect the HbA<sub>1c</sub> levels in Chinese and Malays.

Mean HbA<sub>1c</sub> levels decreased initially from early gestation to a nadir at 21-24 weeks and returned to initial levels at the last trimester. Variable trends had been observed in other studies. Some have reported no change of HbA<sub>1c</sub> with stage of gestation<sup>(9)</sup>, while others have found a fall with increasing gestation<sup>(10, 11)</sup> or a biphasic trend with a nadir level observed at mid-gestation<sup>(12-14)</sup>. These differences had been attributed to difference in methodology and study design. Many of the earlier studies were

cross-sectional in design and used ion-exchange techniques which are sensitive to changes in temperature, acute changes in blood glucose concentrations and interference from minor haemoglobins<sup>(6)</sup>. In our study, HbA<sub>1c</sub> was analysed using affinity chromatography, a method which specifically measures HbA<sub>1c</sub> with minimal interference from labile fractions, pH and temperature changes<sup>(6, 15-17)</sup>. Our results showed a similar trend with those of Morris et al<sup>(12)</sup> who also used the same methodology for analysing HbA<sub>1c</sub> albeit in a longitudinal study. They observed that in their normal pregnant women, HbA<sub>1c</sub> levels decreased progressively from 11-14 weeks, reaching a nadir of 5.2% at 23-26 weeks and returned to baseline concentration (5.6%) by 31-34 weeks. Worth et al<sup>(14)</sup> using 2 methods (a colorimetric and an ion-exchange method), measured HbA<sub>1c</sub> levels sequentially at <12, 17, 24 and 34 weeks and found the nadir at 17 weeks. Phelps et al<sup>(13)</sup> estimating HbA<sub>1c</sub> in a cross-sectional study on normal pregnancy, also observed a biphasic trend with a nadir level at 24 weeks while 1h post-glucose challenge levels followed a similar trend with a nadir level at 4 weeks earlier. They concluded that the changes in HbA<sub>1c</sub> reflected with appropriate displacement in time, the biphasic alterations in mean blood glucose that characterised the sequential changes in glucoregulation during normal pregnancy. In our study, the lower mean random plasma glucose level at 17-20 weeks might probably reflect the nadir level of mean HbA<sub>1c</sub> 4 weeks later. However, 1h post-glucose challenge responses in our study did not show an obvious trend.

In cases where a GTT was done (221 cases), the variation of HbA<sub>1c</sub> with gestation showed a similar trend with nadir at 21-24 weeks but a slight peak at 33-36 weeks (Fig 2). The difference towards the later stage of gestation may be due to the difference in case number for the gestation groups. The trend of glucose levels in 0h, 1h or 2h GTT did not appear to follow closely the trend of HbA<sub>1c</sub> levels. However, the lower levels of 0h GTT at 17-20 and 21-24 weeks gestation periods may have contributed to the nadir level of HbA<sub>1c</sub> at 17-20 weeks.

Theoretically HbA<sub>1c</sub> values could be used as a simple and convenient screening test for gestational diabetes. However, investigators have not been able to find a clear cut-off point that will differentiate between normal and abnormal glucose tolerance in pregnancy. This overlap and the wide variation in the range of HbA<sub>1c</sub> values between the 2 groups does not make HbA<sub>1c</sub> an accurate screening test for gestational diabetes. HbA<sub>1c</sub> may be used as an additional parameter for assessment of glucose control and a lower range (compared to non-pregnant women) shown here maybe used as a guideline for good control.

## ACKNOWLEDGEMENTS

The authors would like to thank staff nurses, Messrs YN Ong, Lye YC, Tham WC, Chan SC, Leong FY and Ng KF and technicians, Messrs LG Tan and SC Kua for their technical assistance and Associate Professor Lun Kwok Chye for advice on statistical analysis and the National University of Singapore for their generous support of this study.

## REFERENCES

- Gabby KH. Glycosylated haemoglobin and diabetes control (Editorial). *N Engl J Med* 1976; 295: 443-4.
- Kennedy AL, Merimee TJ. Glycosylated serum protein and hemoglobin A1 levels to measure control of glycemia. *Ann Intern Med* 1981; 95: 56-8.
- Larsen ML, Horder M, Mogensen EF. Effect of long-term monitoring of glycosylated haemoglobin levels in insulin-dependent diabetes mellitus. *N Engl J Med* 1990; 323: 1021-5.
- Chua S, Thai AC, Kek LP, Yeoh SC, Ratnam SS. The glucose challenge test: A screening test for gestational diabetes mellitus.

- Singapore Med J 1993; 34: 303-5.
5. Bondar RJL, Mead DC. Evaluation of glucose-6-phosphate dehydrogenase for leuconostoc mesenteroides in the hexokinase method for determining glucose in serum. Clin Chem 1974; 20: 586-90.
  6. Klenk DC, Hermanson GT, Krohn JI, Fujimoto EK, Mallia AK, Smith PK, et al. Determination of glycosylated hemoglobin by affinity chromatography: comparison with colorimetric and ion-exchange methods and effects of common interferences. Clin Chem 1982; 28: 2088-94.
  7. World Health Organization Study Group Report. Diabetes Mellitus. WHO Tech Rep Ser 1985; 727: 9-20.
  8. Liew-Loke DFM, Kek LP, Thai AC, Ng CSA, Ratnam SS. Haemoglobin A<sub>1c</sub> levels in non-diabetic pregnant and non-pregnant women. Sing J Obstet Gynaecol 1988; 19: 18-21.
  9. O' Shaughnessy R, Russ RJ, Zuspan F. Glycosylated hemoglobins and diabetes mellitus in pregnancy. Am J Obstet Gynecol 1979; 135: 783-90.
  10. Hanson U, Hagenfeldt L, Hagenfeldt K. Glycosylated hemoglobins in normal pregnancy: sequential change and relation to birth weight. Obstet Gynecol 1983; 62: 741-4.
  11. Widness JA, Schwartz HC, Kahn CB, Oh W, Schwartz R. Glycohemoglobin determinations in diabetic pregnancy: a sequential study. Am J Obstet Gynecol 1980; 136: 1024-9.
  12. Morris MN, Grandis AS, Litton JC. Longitudinal assessment of glycosylated blood protein concentrations in normal pregnancy and gestational diabetes. Diab Care 1986; 9: 107-10.
  13. Phelps RL, Honig G, Green D, Metzger BE, Frederiksen MC, Freinkel N. Biphasic changes in hemoglobin A<sub>1c</sub> concentrations during normal human pregnancy. Am J Obstet Gynecol 1983; 147: 651-3.
  14. Worth R, Potter JM, Drury J, Fraser RB, Cullen Dr. Glycosylated haemoglobin in normal pregnancy: a longitudinal study with two independent methods. Diabetologia 1985; 28: 76-7.
  15. Gould BJ, Hall PM, Cook JGH. Measurement of glycosylated hemoglobins using an affinity chromatography method. Clin Chim Acta 1982; 125: 41-8.
  16. Talwar D, Barr BB, Kesson CM, Robb DA. Determination of glycosylated adult and fetal hemoglobins by affinity chromatography. Clin Chim Acta 1983; 128: 61-7.
  17. Yue DK, McLennan S, Church DB, Turtle JR. The measurement of glycosylated hemoglobin in man and animals by aminophenyl boronic acid affinity chromatography. Diabetes 1982; 31: 701-5.

## 3rd Asian Congress on Oral & Maxillofacial Surgery

organised by the Asian Association  
of Oral & Maxillofacial Surgeons

28 – 31 March 1996

Venue: Hilton Hotel  
Kuching, Sarawak  
Malaysia

For further information, please contact:

Dr N Ravindranathan  
Organising Chairman  
c/o Agas-Pan Asian Sdn Bhd  
2nd Floor, Unit R206  
Sarawak Plaza  
Jalan Tunku Abdul Rahman  
P O Box 3208, 93762 Kuching  
Sarawak, Malaysia  
Tel: (6082) 428969/429754  
Fax: (6082) 419754