

A variation of the phrenic nerve: case report and review

Prakash, Prabhu L V, Madhyastha S, Singh G

ABSTRACT

During routine dissection in the department of anatomy, the following anatomical variations of the phrenic nerve were observed on the right side of the neck of a 30-year-old male cadaver. The phrenic nerve, in its early course close to its origin, gave a communicating branch to the C5 root of the brachial plexus. At the level of the root of neck just before entering the thorax, the phrenic nerve was located anterior to the subclavian vein. This unique case of phrenic nerve variation gains tremendous importance in the context of subclavian vein cannulation, implanted venous access portals, and supraclavicular nerve block for regional anaesthesia.

Keywords: anatomical variations, brachial plexus, neck anatomy, phrenic nerve variations.

Singapore Med J 2007;48(12):1156-1157

INTRODUCTION

Variations of the phrenic nerve have been observed and reported in the past.⁽¹⁻³⁾ Right subclavian vein cannulation is a commonly-performed procedure for vascular access for haemodialysis, and the phrenic nerve palsy manifesting as hemidiaphragmatic paralysis is an important complication associated with it.⁽⁴⁻⁸⁾ Right phrenic nerve paralysis is an acknowledged immediate and late complication of implanted venous access portals.⁽⁹⁾ Bigeleisen reported a case of simultaneous diaphragmatic and brachial plexus stimulation, followed by a successful nerve block, and demonstrated the necessity of a thorough knowledge of anatomical variations and standard anatomy, for the safe and efficient practice for regional anaesthesia.⁽¹⁰⁾

CASE REPORT

The following important variations were observed only on the right side during routine dissection in a 30-year-old male cadaver. The phrenic nerve, in its early course close to its origin, gave a communicating branch to the C5 root of the brachial plexus (Fig. 1).



Fig. 1 Photograph shows the variations of the phrenic nerve. AC: ansa cervicalis; CB: communicating branch to brachial plexus; C5: C5 root of brachial plexus; PN: phrenic nerve; IJV: Internal jugular vein; SCV: subclavian vein

Department of Anatomy, Centre for Basic Sciences, Kasturba Medical College, Bejei, Mangalore 575004, Karnataka, India

Prakash, MBBS, MD Assistant Professor

Prabhu LV, MBBS, MS Professor and Head

Madhyastha S, MSc, PhD Associate Professor

Department of Anatomy, Institute of Medical Sciences, Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India

Gajendra Singh, MBBS, MS Professor and Director

Correspondence to: Dr Prakash Department of Anatomy, Vyhedi Institute of Medical Sciences & Research Centre, 82 EPIP Area, Whitefield, Bangalore 560066, Karnataka, India Tel: (91) 9480 229 670 Fax: (91) 8028 416 199 Email: prakashrinku@rediffmail.com

The phrenic nerve at the level of the root of the neck, just before entering the thorax, was located in front of the subclavian vein (Fig. 1). It is usually located posteriorly, in between subclavian vein and artery. This variation of phrenic nerve makes it highly vulnerable to injury during subclavian catheterisation for vascular access.

DISCUSSION

The accessory phrenic nerve proceeds caudally and joins distally with phrenic nerve, and is present in up to 75% of cadavers.⁽¹⁰⁾ On the contrary, in this particular case, the variation is not an accessory phrenic nerve; as the communicating branch from the phrenic nerve to the C5 root of the brachial plexus proceeded distally and laterally in caudal direction and joined the C5 root of the brachial plexus. Hollinshead described that the entire phrenic nerve, in ten out of 138 (7.25%) sides of cadavers studied, passed anterior to subclavian vein as it leaves the neck,⁽¹⁾ whereas in the present case, this particular variation of phrenic nerve was accompanied further by a communicating branch to the C5 root of the brachial plexus.

Bigeleisen reported that more distal blocks, such as a low interscalene or supraclavicular block, may also give rise to a partial phrenic nerve block, even if the phrenic nerve anatomy is standard.⁽¹⁰⁾ In the context of the present case, this will lead to complete phrenic nerve block manifesting as right diaphragmatic paralysis. The guidance of the developing axons is regulated by expression of chemoattractants and chemorepulsants, in a highly-coordinated site specific fashion. Any alterations in signalling between mesenchymal cells and neuronal growth cones can lead to significant variations, which probably in the present case, resulted in phrenic nerve communication with the C5 root of the brachial plexus, along with subclavian vein located posterior to the phrenic nerve. Once formed, any developmental differences would persist postnatally.⁽¹¹⁾

REFERENCES

- Hollinshead WH. Anatomy for Surgeons. The Head and Neck. 3rd ed. New York: Harper and Row Publishers, 1982.
- Anson BJ, McVay CB. Surgical Anatomy. 5th ed. Philadelphia: WB Saunders, 1971.
- Bergman RA, Thompson SA, Afifi AK, Saddeh FA. Cependium of Human Anatomical Variation. Baltimore: Urban and Schwarzenburg, 1988.
- Islek I, Akpolat T, Danaci M. Phrenic nerve palsy caused by subclavian vein catheterization. Nephrol Dial Transplant, 1998; 13:1023-5.
- Porzionato A, Montisci M, Manani G. Brachial plexus injury following subclavian vein catheterization: a case report. J Clin Anesth 2003; 15:582-6.
- Akata T, Noda Y, Nagata T, Noda E, Kandabashi T. Hemidiaphragmatic paralysis following subclavian vein catheterization. Acta Anaesthesiol Scand 1997; 41:1223-5.
- D'Netto MA, Bender J, Brown RT, Herson VC. Unilateral diaphragmatic palsy in association with a subclavian vein thrombus in a very-low-birth weight infant. Am J Perinatol 2001; 18:459-64.
- Aggarwal S, Hari P, Bagga A, Mehta SN. Phrenic nerve palsy: a rare complication of indwelling subclavian vein catheter. Pediatr Nephrol 2000; 14:203-4.
- Reeves JE Jr, Anderson WF. Permanent paralysis of the right phrenic nerve. Ann Intern Med 2002; 137:551-2.
- Bigeleisen PE. Anatomical variations of the phrenic nerve and its clinical implication for supraclavicular block. Br J Anaesth 2003; 91:916-7.
- Sanes DH, Reh TA, Harris WA. Development of the Nervous System. London: Academic Press, 2000.

So many choices, too little time.
Tired of channel surfing?



Singapore Medical Journal. The perfect channel for your publication.

The voice of academic medicine in Singapore and Southeast Asia since 1960.

Tel: 65 6223 1264 Fax: 65 6224 7827 Email: smj@sma.org.sg

www.sma.org.sg/smj